

Status of the $B_{(s)}^0$ - $\bar{B}_{(s)}^0$ mixing from QCD spectral sum Rules ¹

Stephan Narison

Laboratoire de Physique Mathématique
Université de Montpellier II
Place Eugène Bataillon
34095 - Montpellier Cedex 05, France
Email: qcd@lpm.univ-montp2.fr

Abstract

In this talk, I present new results [1] obtained from QCD spectral sum rules (QSSR), on the bag constant parameters entering in the analysis of the $B_{(s)}^0$ - $\bar{B}_{(s)}^0$ mass-differences. Taking the average of the results from the Laplace and moment sum rules, one obtains to order α_s : $f_B \sqrt{\hat{B}_B} \simeq (228 \pm 61)$ MeV, $f_{B_s} \sqrt{\hat{B}_{B_s}} / f_B \sqrt{\hat{B}_B} \simeq 1.18 \pm 0.03$, in units where $f_\pi = 130.7$ MeV. Combined with the experimental data on the mass-differences $\Delta M_{d,s}$, one obtains the constraint on the CKM weak mixing angle: $|V_{ts}/V_{td}|^2 \geq 20.2(1.3)$. Alternatively, using the weak mixing angle from the analysis of the unitarity triangle and the data on ΔM_d , one predicts $\Delta M_s = 18.3(2.1) \text{ ps}^{-1}$ in agreement with the present experimental lower bound and within the reach of Tevatron 2.

1 Introduction

$B_{(s)}^0$ and $\bar{B}_{(s)}^0$ are not eigenstates of the weak hamiltonian, such that their oscillation frequency is governed by their mass-difference ΔM_q . The measurement by the UA1 collaboration [2] of a large value of ΔM_d was the *first* indication of a heavy top-quark mass. In the SM, the mass-difference is approximately given by [3]:

$$\Delta M_q \simeq \frac{G_F^2}{4\pi^2} M_W^2 |V_{tq} V_{tb}^*|^2 S_0 \left(\frac{m_t^2}{M_W^2} \right) \eta_B C_B(\nu) \frac{1}{2M_{B_q}} \langle \bar{B}_q^0 | \mathcal{O}_q(\nu) | B_q^0 \rangle \quad (1)$$

where the $\Delta B = 2$ local operator $\mathcal{O}_q$ is defined as:

$$\mathcal{O}_q(x) \equiv (\bar{b} \gamma_\mu L q)(\bar{b} \gamma_\mu L q) , \quad (2)$$

¹Talk presented at the 1st High-Energy Physics Madagascar International Conference Series (HEP-MAD'01), 27th sept-5th Oct. 2001, Antananarivo.

with: $L \equiv (1 - \gamma_5)/2$ and $q \equiv d, s$; S_0 , η_B and $C_B(\nu)$ are short distance quantities and Wilson coefficients which are calculable perturbatively, while the matrix element $\langle \bar{B}_q^0 | \mathcal{O}_q | B_q^0 \rangle$ requires non-perturbative QCD calculations, and is usually parametrized for $SU(N)_c$ colours as:

$$\langle \bar{B}_q^0 | \mathcal{O}_q | B_q^0 \rangle = N_c \left(1 + \frac{1}{N_c} \right) f_{B_q}^2 M_{B_q}^2 B_{B_q} . \quad (3)$$

f_{B_q} is the B_q decay constant normalized as $f_\pi = 92.4$ MeV, and B_{B_q} is the so-called bag parameter which is $B_{B_q} \simeq 1$ if one uses a vacuum saturation of the matrix element. From Eq. (1), it is clear that the measurement of ΔM_d provides the one of the CKM mixing angle $|V_{td}|$ if one uses $|V_{tb}| \simeq 1$. One can also extract this quantity from the ratio:

$$\frac{\Delta M_s}{\Delta M_d} = \left| \frac{V_{ts}}{V_{td}} \right|^2 \frac{M_{B_d}}{M_{B_s}} \frac{\langle \bar{B}_s^0 | \mathcal{O}_s | B_s^0 \rangle}{\langle \bar{B}_d^0 | \mathcal{O}_d | B_d^0 \rangle} \equiv \left| \frac{V_{ts}}{V_{td}} \right|^2 \frac{M_{B_d}}{M_{B_s}} \xi^2 , \quad (4)$$

since in the SM with three generations and unitarity constraints, $|V_{ts}| \simeq |V_{cb}|$. Here:

$$\xi \equiv \sqrt{\frac{g_s}{g_d}} \equiv \frac{f_{B_s} \sqrt{B_{B_s}}}{f_B \sqrt{B_B}} . \quad (5)$$

The great advantage of Eq. (4) compared with the former relation in Eq. (1) is that in the ratio, different systematics in the evaluation of the matrix element tends to cancel out, thus providing a more accurate prediction. However, unlike $\Delta M_d = 0.473(17) \text{ ps}^{-1}$, which is measured with a good precision [4], the determination of ΔM_s is an experimental challenge due to the rapid oscillation of the B_s^0 - $\bar{B}_s^0$ system. At present, only a lower bound of 13.1 ps^{-1} is available at the 95% CL from experiments [4], but this bound already provides a strong constraint on $|V_{td}|$.

2 Two-point function sum rule

Ref. [5] has extended the analysis of the K^0 - $\bar{K}^0$ systems of [6], using two-point correlator of the four-quark operators into the analysis of the quantity $f_B \sqrt{B_B}$ which governs the B^0 - $\bar{B}^0$ mass difference. The two-point correlator defined as:

$$\psi_H(q^2) \equiv i \int d^4x e^{iqx} \langle 0 | \mathcal{T} \mathcal{O}_q(x) (\mathcal{O}_q(0))^\dagger | 0 \rangle , \quad (6)$$

is built from the $\Delta B = 2$ weak operator defined in Eq. (2). The corresponding Laplace (resp. moment) sum rules are:

$$\mathcal{L}(\tau) = \int_{4M_B^2}^{\infty} dt e^{-t\tau} \text{Im} \psi_H(t) , \quad \mathcal{M}_n = \int_{4M_B^2}^{\infty} dt t^n \text{Im} \psi_H(t) , \quad (7)$$

The two-point function approach is very convenient due to its simple analytic properties which is not the case of approach based on three-point functions². However, it involves non-trivial QCD calculations which become technically complicated when one includes the contributions of radiative corrections due to non-factorizable diagrams. These perturbative radiative corrections due to *factorizable* and *non-factorizable* diagrams have been already computed in [8] (referred as NP), where it has been found that the factorizable corrections are large while the non-factorizable ones are negligibly small. NP analysis has confirmed the estimate in [5] from lowest order calculations, where under some assumptions on the contributions of higher mass resonances to the spectral function, the value of the bag parameter B_B has been found to be:

$$B_{B_d}(4m_b^2) \simeq (1 \pm 0.15) . \quad (8)$$

²For detailed criticisms, see [7].

This value is comparable with the one $B_{B_d} = 1$ from the vacuum saturation estimate, which is expected to be a quite good approximation due to the relative high-scale of the B -meson mass. Equivalently, the corresponding RGI quantity is:

$$\hat{B}_{B_d} \simeq (1.5 \pm 0.2) \quad (9)$$

where we have used the relation:

$$B_{B_q}(\nu) = \hat{B}_{B_q} \alpha_s^{-\frac{\gamma_0}{\beta_1}} \left\{ 1 - \left(\frac{5165}{12696} \right) \left(\frac{\alpha_s}{\pi} \right) \right\}, \quad (10)$$

with $\gamma_0 = 1$ is the anomalous dimension of the operator $\mathcal{O}_q$ and $\beta_1 = -23/6$ for 5 flavours. The NLO corrections have been obtained in the $\overline{MS}$ scheme [3]. We have also used, to this order, the value [9, 7]:

$$\bar{m}_b(m_b) = (4.24 \pm 0.06) \text{ GeV}, \quad (11)$$

and $\Lambda_5 = (250 \pm 50) \text{ MeV}$ [10]. In a forthcoming paper [1], we study (*for the first time*), from the QCD spectral sum rules (QSSR) method, the $SU(3)$ breaking effects on the ratio: ξ defined previously in Eq. (5), where a similar analysis of the ratios of the decay constants has given the values [11]:

$$\frac{f_{D_s}}{f_D} \simeq 1.15 \pm 0.04, \quad \frac{f_{B_s}}{f_B} \simeq 1.16 \pm 0.04. \quad (12)$$

We also improve the previous result on B_{B_d} by the inclusion of the $B^* - B^*$ resonances into the spectral function.

3 Results and implications on $|V_{ts}/V_{td}|^2$ and ΔM_s

We deduce by taking the average from the moments and Laplace sum rules results [1]:

$$\xi \equiv \frac{f_{B_s} \sqrt{B_{B_s}}}{f_B \sqrt{B_B}} \simeq 1.18 \pm 0.03, \quad f_B \sqrt{B_B} \simeq (228 \pm 61) \text{ MeV}, \quad (13)$$

in units where $f_\pi = 130.7 \text{ MeV}$. As expected, we have smaller errors for the ratio ξ due to the cancellation of the systematics, while for $f_B \sqrt{B_B}$, the error comes mainly and equally from the pole mass M_b and the truncation of the PT series where we have estimated the strength of the α_s^2 contribution assuming a geometric growth of the PT coefficients. These results can be compared with different lattice and phenomenological determinations given in [12, 13]. By comparing the ratio with the one of f_{B_s}/f_{B_d} in Eq. (12)³, one can conclude (to a good approximation) that:

$$\hat{B}_{B_s} \simeq \hat{B}_{B_d} \simeq (1.41 \pm 0.33) \implies B_{B_{d,s}}(4m_b^2) \simeq (0.94 \pm 0.22), \quad (14)$$

indicating a negligible $SU(3)$ breaking for the bag parameter. For a consistency, we have used the estimate to order α_s [15]:

$$f_B \simeq (1.47 \pm 0.10) f_\pi, \quad (15)$$

and we have assumed that the error from f_B compensates the one in Eq. (13). The result is in excellent agreement with the previous result of [8] in Eqs (8) and (9). Using the experimental values:

$$\Delta M_d = 0.472(17) \text{ ps}^{-1}, \quad \Delta M_s \geq 13.1 \text{ ps}^{-1} \quad (95\% \text{ CL}), \quad (16)$$

one can deduce from Eq. (4):

$$\rho_{sd} \equiv \left| \frac{V_{ts}}{V_{td}} \right|^2 \geq 20.2(1.3). \quad (17)$$

³One can notice that similar strengths of the $SU(3)$ breakings have been obtained for the $B \rightarrow K^* \gamma$ and $B \rightarrow K \ell \nu$ form factors [14].

Alternatively, using: $\rho_{sd} \simeq 28.4(2.9)$ obtained by using the Wolfenstein parameters determined in [12], we deduce:

$$\Delta M_s \simeq 18.4(2.1) \text{ } ps^{-1} \text{ ,} \quad (18)$$

in good agreement with the present experimental lower bound and within the reach of Tevatron run 2 experiments.

4 Conclusions

We have applied QCD spectral sum rules for extracting (*for the first time*) the $SU(3)$ breaking parameter in Eq. (13). The phenomenological consequences of our results for the $B_{d,s}^0 - \bar{B}_{d,s}^0$ mass-differences and CKM mixing angle have been discussed. An extension of this work to the study of the $B_{s,d}^0 - \bar{B}_{s,d}^0$ width difference is in progress [16].

Acknowledgements

It is a pleasure to thank my collaborators Kaoru Hagiwara and Daisuke Nomura.

References

- [1] D. Nomura, K. Hagiwara and S. Narison (to appear).
- [2] C. Albajar et al., UA1 collaboration, *Phys. Lett. B* **186**, 237, 247 (1987).
- [3] G. Buchalla, A.J. Buras and M.E. Lautenbacher, *Rev. Mod. Phys.* **68**, 1125 (1996).
- [4] PDG, D.E. Groom et al., *EPJ C* **15**, 1 (2000) and 2001 off-year partial update for the 2002 edition, <http://pdg.lplbl.gov/>.
- [5] A. Pich, *Phys. Lett. B* **206**, 322 (1988).
- [6] B. Guberina et al., *Phys. Lett. B* **128**, 269 (1983) ; A. Pich and E. de Rafael, *Phys. Lett. B* **158**, 477 (1985); *Nucl. Phys. B* **358**, 311 (1991); S. Narison, *Phys. Lett. B* **351**, 369 (1995).
- [7] For a review and references to original works, see e.g., S. Narison, *QCD as a Theory of Hadrons*, Cambridge Press, Monograph Series 2002 (to appear); *QCD spectral sum rules*, *World Sci. Lect. Notes Phys.* **26**, 1 (1989).; *Phys. Rep.* **84**, 263 (1982).
- [8] S. Narison and A. Pivovarov, *Phys. Lett. B* **327** 341 (1994).
- [9] S. Narison, *Phys. Lett. B* **341**, 73 (1994); *Phys. Lett. B* **520** 1152001.
- [10] S. Bethke, *Nucl. Phys. (Proc. Suppl.) B*, A **54**, 314 (1997); hep-ex/0004201.
- [11] S. Narison, *Phys. Lett. B* **320**, 369 (1994).
- [12] For a recent compilation, see e.g. M.Ciuchini et al., hep-ph/0012308.
- [13] C. Bernard, *Nucl. Phys. (Proc. Suppl.) B* **94**, 159 (2001) and references therein.
- [14] S. Narison, *Phys. Lett. B* **327**, 354 (1994); B **337** 163 (1994); B **283**, 384 (1992).
- [15] S. Narison, *Nucl. Phys. (Proc. Suppl.) B* **74**, 304 (1999).
- [16] D. Nomura, K. Hagiwara and S. Narison (to appear).